

National College Health Assessment Comparison

Kent State 2003 responses are from the American College Health Association. National College Health Assessment: Kent State University Executive Summary Fall 2003. Baltimore: American College Health Association; 2003. (Sample size (n) = 655))

Kent State 2004 responses are from the American College Health Association. National College Health Assessment: Reference Group Executive Summary Fall 2004. Baltimore: American College Health Association; 2004. (Sample size (n) = 601))

Kent State 2008 responses are from the American College Health Association. National College Health Assessment: Kent State University Executive Summary Spring 2008. Baltimore: American College Health Association; 2008. (Sample size (n) = 427))

Reference group 2007 responses are from the American College Health Association. National College Health Assessment: Reference Group Executive Summary Fall 2007. Baltimore: American College Health Association; 2007. (Sample size (n) = 20,507))

> Data analysis performed by Donna J. Coto, Coordinator of Health Services Information

Kent State University – Health Services **All numbers are percentage of the total**

A. General Health of College Students (Alphabetical)

Within the last 12 months, college students reported experiencing:

| | KSU | KSU | KSU | Reference |
|-----------------------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| Allergy Problems | 53.2 | 60.4 | 48.6 | 51.0 |
| Anorexia | 2.2 | 1.5 | 2.8 | 1.8 |
| Anxiety Disorder | 16.4 | 16.4 | 18.2 | 13.1 |
| Asthma | 9.1 | 13.0 | 12.8 | 12.1 |
| Bulimia | 4.1 | 1.3 | 3.1 | 2.0 |
| Chronic Fatigue Syndrome | 5.4 | 4.4 | 4.9 | 3.4 |
| Depression | 25.0 | 21.0 | 20.6 | 18.9 |
| Diabetes | 2.0 | 2.0 | 1.7 | 1.5 |
| Endometriosis | 2.5 | 2.5 | 1.2 | 1.3 |
| Genital Herpes | 1.7 | 0.7 | 2.1 | 1.0 |
| Genital Warts/HPV | 2.2 | 2.5 | 3.1 | 2.3 |
| Hepatitis B or C | 0.8 | 0.7 | 0.7 | 0.5 |
| High Blood Pressure | 7.3 | 5.9 | 5.5 | 6.3 |
| High Cholesterol | 4.6 | 3.9 | 2.8 | 4.4 |
| HIV Infection | 0.3 | 0.7 | 0.5 | 0.3 |
| Repetitive Stress Injury | 8.9 | 8.4 | 6.1 | 7.4 |
| Seasonal Affective Disorder | 10.4 | 8.9 | 13.2 | 6.8 |
| Substance Abuse Problem | 3.4 | 5.2 | 4.5 | 3.6 |
| Back Pain | 55.7 | 58.2 | 55.0 | 49.4 |
| Broken Bone/Fracture | 6.0 | 5.9 | 5.4 | 5.5 |
| Bronchitis | 8.9 | 10.8 | 10.6 | 7.9 |
| Chlamydia | 1.2 | 0.8 | 1.4 | 0.9 |
| Ear Infection | 13.7 | 14.0 | 10.2 | 11.5 |
| Gonorrhea | 0.5 | 0.3 | 0.5 | 0.3 |
| Mononucleosis | 2.2 | 3.9 | 4.0 | 2.2 |
| Pelvic Inflammatory Disease | 0.6 | 0.7 | 0.5 | 0.4 |
| Sinus Infection | 42.8 | 43.6 | 42.5 | 32.6 |
| Strep Throat | 16.0 | 16.1 | 16.7 | 15.5 |
| Tuberculosis | 0.2 | 0.2 | 0.0 | 0.4 |

B. Preventative Health

Preventative health practices among college students:

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|--------------------------------|-------------|-------------|-------------|------------------|
| Hepatitis B Vaccination | 54.1 | 55.2 | 58.4 | 68.7 |
| Meningitis Vaccination | 39.2 | 38.6 | 53.5 | 52.8 |
| Varicella Vaccination | 48.4 | 49.6 | 48.9 | 52.7 |
| MMR Vaccination | 85.9 | 83.9 | 84.2 | 77.1 |
| Influenza Vaccination | 20.0 | 19.4 | 23.9 | 29.2 |
| Dental Exam in Last Year | 76.6 | 74.9 | 77.7 | 76.3 |
| Testicular Self-Exam in Last | 35.7 | 34.0 | 47.1 | 42.1 |
| Month | | | | |
| Breast Self-Exam in Last Month | 48.7 | 50.4 | 46.8 | 46.5 |
| Gynecological Exam in Last | 65.5 | 63.2 | 64.4 | 55.8 |
| Year | | | | |
| Blood Pressure Checked in Last | 91.3 | 91.4 | 88.9 | 87.9 |
| 2 Years | | | | |
| Cholesterol Checked in Last 5 | 41.7 | 43.1 | 40.7 | 43.8 |
| Years | | | | |
| Sunscreen Daily Use | 14.9 | 14.2 | 17.6 | 18.8 |

C. Academic Impacts (Alphabetical)

Within the last 12 months percentage of college students reported the following factors affecting their individual academic performance:

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|--|-------------|-------------|-------------|------------------|
| Alcohol Use | 7.4 | 7.4 | 8.7 | 6.2 |
| Allergies | 6.2 | 5.9 | 5.4 | 5.2 |
| Assault (Physical) | 1.1 | 0.7 | 0.9 | 0.7 |
| Assault (Sexual) | 1.2 | 0.3 | 1.2 | 0.9 |
| Attention Deficit Disorder | 6.8 | 8.8 | 8.8 | 7.0 |
| Cold/Flu/Sore Throat | 26.7 | 26.1 | 27.2 | 24.8 |
| Concern for Friend/Family Member | 20.1 | 20.9 | 17.5 | 18.1 |
| Chronic Illness | 2.6 | 5.6 | 4.3 | 3.2 |
| Chronic Pain | 4.2 | 3.9 | 4.3 | 3.8 |
| Death of a Friend/Family Member | 9.0 | 10.3 | 9.2 | 9.8 |
| Depression/Anxiety Disorder/SAD | 20.0 | 18.2 | 19.9 | 15.5 |
| Drug Use | 2.5 | 2.0 | 2.9 | 2.1 |
| Eating Disorder/Problem | 2.2 | 0.7 | 1.2 | 1.3 |
| HIV Infection | 0.3 | 0.2 | 0.5 | 0.2 |
| Injury | 3.8 | 3.1 | 3.8 | 4.1 |
| Internet Use/Computer Games | 11.7 | 13.1 | 14.7 | 15.1 |
| Learning Disability | 3.7 | 3.9 | 4.5 | 4.0 |
| Mononucleosis | 1.5 | 2.2 | 1.4 | 1.4 |
| Pregnancy (Yours or Partner's) | 1.1 | 0.7 | 1.4 | 1.4 |
| Relationship Difficulty | 17.2 | 17.0 | 18.4 | 15.5 |
| Sexually Transmitted Disease | 0.8 | 0.0 | 0.7 | 0.5 |
| Sinus/Ear/Bronchitis/Strep Infections | 11.0 | 11.0 | 8.5 | 9.4 |
| Sleep Difficulties | 26.5 | 30.3 | 27.2 | 25.4 |
| Stress | 37.2 | 34.0 | 34.4 | 32.9 |

D. Violence

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|---|-------------|-------------|-------------|---------------------|
| A physical fight | 6.8 | 8.5 | 7.1 | 8.3 |
| Being physically assaulted (non-sexual) | 4.5 | 4.5 | 5.0 | 4.3 |
| Verbal threats for sex against their will | 4.6 | 4.3 | 4.7 | 3.5 |
| Sexual touching against their will | 12.5 | 8.4 | 6.8 | 6.9 |
| Attempted penetration against their will | 4.6 | 4.2 | 3.5 | 2.8 |
| Penetration against their will | 2.4 | 1.8 | 1.7 | 1.7 |
| An emotionally abusive relationship | 18.8 | 15.5 | 17.9 | 14.1 |
| A physically abusive relationship | 2.6 | 2.7 | 2.1 | 2.4 |
| A sexually abusive relationship | 1.4 | 2.0 | 1.6 | 2.0 |

Within the last 12 months, college students reported experiencing:

E. Alcohol, Tobacco, and Other Drug Use

Thirty day prevalence – substance use reported by college students:

| Alcohol | | | | |
|------------------------------|------|------|------|-----------|
| | KSU | KSU | KSU | Reference |
| | 2003 | 2004 | 2008 | (2007) |
| Never used | 16.0 | 15.2 | 15.0 | 22.8 |
| Used, but not in the last 30 | 15.2 | 16.8 | 18.5 | 15.4 |
| days | | | | |
| Used 1-9 days | 53.1 | 52.9 | 51.6 | 47.7 |
| Used 10-29 days | 14.9 | 14.9 | 13.6 | 13.4 |
| Used all 30 days | 0.8 | 0.2 | 1.2 | 0.7 |

Cigarette

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------------------------|-------------|-------------|-------------|------------------|
| Never used | 53.1 | 56.1 | 57.1 | 65.6 |
| Used, but not in the last 30 | 17.5 | 19.2 | 21.5 | 15.5 |
| days | | | | |
| Used 1-9 days | 11.7 | 11.7 | 10.1 | 8.9 |
| Used 10-29 days | 7.4 | 6.5 | 3.7 | 4.1 |
| Used all 30 days | 10.4 | 6.5 | 7.5 | 5.9 |

Marijuana

| | KSU | KSU | KSU | Reference |
|------------------------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| Never used | 67.5 | 66.8 | 64.1 | 68.8 |
| Used, but not in the last 30 | 19.3 | 21.7 | 21.8 | 18.5 |
| days | | | | |
| Used 1-9 days | 8.7 | 8.2 | 9.2 | 8.4 |
| Used 10-29 days | 3.5 | 3.2 | 3.3 | 3.0 |
| Used all 30 days | 0.9 | 0.2 | 1.6 | 1.2 |

Reported number of times college students consumed five or more drinks in a sitting within the last two weeks:

| | KSU | KSU | KSU | Reference |
|-----------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| None | 63.9 | 61.8 | 60.6 | 66.5 |
| 1-2 times | 22.9 | 26.0 | 24.2 | 20.9 |
| 3-5 times | 10.1 | 8.8 | 11.3 | 9.8 |
| 6 or more times | 3.1 | 3.3 | 4.0 | 2.8 |

When at a party or going out (usually or always)...

| vinen at a party of going out (a | KSU | KSU | KSU | Reference |
|----------------------------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| Alternate non-alcoholic with | 30.2 | 26.8 | 27.2 | 33.6 |
| alcoholic beverages | | | | |
| Determine in advance not to | 41.3 | 38.3 | 36.8 | 43.4 |
| exceed a set number of drinks | | | | |
| Choose not to drink alcohol | 30.9 | 30.9 | 28.7 | 29.5 |
| Use a designated driver | 79.8 | 77.2 | 85.3 | 80.6 |
| Eat before and/or during | 76.0 | 77.3 | 82.4 | 81.2 |
| drinking | | | | |
| Have a friend let you know | 27.9 | 27.9 | 26.1 | 31.4 |
| when you've had enough | | | | |
| Keep track of how many drinks | 67.0 | 67.9 | 64.0 | 66.4 |
| being consumed | | | | |
| Pace drinks to one or fewer an | 34.9 | 29.6 | 32.9 | 34.0 |
| hour | | | | |
| Avoid drinking games | 51.4 | 46.7 | 37.7 | 42.1 |
| Drink an alcoholic look-alike | 7.9 | 8.7 | 6.5 | 8.5 |

*Students who responded "N/A don't drink" were excluded from this analysis

| in the last 12 months as a rest | KSU | KSU | KSU | Reference |
|---------------------------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| Being physically injured | 15.7 | 18.9 | 20.6 | 15.3 |
| Physically injuring another | 3.8 | 3.0 | 5.7 | 4.1 |
| person | | | | |
| Being involved in a fight | 4.2 | 6.6 | 6.3 | 6.5 |
| Doing something they later | 31.5 | 32.3 | 34.0 | 31.1 |
| regretted | | | | |
| Forgetting where they were | 21.8 | 29.7 | 30.1 | 26.2 |
| or what they had done | | | | |
| Having someone use force or | 2.3 | 1.5 | 1.1 | 1.5 |
| threat of force to have sex | | | | |
| with them | | | | |
| Having unprotected sex | 15.1 | 14.7 | 18.7 | 14.2 |

College students who drank alcohol reported the following consequences occurring in the last 12 months as a result of their own drinking:

*Students who responded "N/A don't drink" were excluded from this analysis

F. Sexual Behavior

College students reported the following within the last 12 months:

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------------------------|-------------|-------------|-------------|------------------|
| Having had no sexual partner | 24.4 | 24.5 | 20.9 | 31.4 |
| Having had 1 sexual partner | 50.8 | 51.1 | 52.0 | 45.3 |
| Having had 2 sexual partners | 11.0 | 11.7 | 11.8 | 10.1 |
| Having had 3 sexual partners | 6.1 | 4.2 | 7.5 | 5.3 |
| Having had 4 or more sexual | 7.7 | 8.5 | 7.8 | 7.9 |
| partners | | | | |

Oral sex within the past 30 days

| | KSU | KSU | KSU | Reference |
|-----------------------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| Never did this activity | 20.8 | 21.9 | 17.9 | 31.3 |
| Have not done this activity | 27.5 | 27.8 | 29.1 | 26.0 |
| during last 30 | | | | |
| Did this one or more times | 51.7 | 50.3 | 53.0 | 42.7 |

Vaginal sex within the past 30 days

| | KSU | KSU | KSU | Reference |
|-----------------------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| Never did this activity | 25.5 | 26.5 | 22.0 | 31.9 |
| Have not done this activity | 21.0 | 22.2 | 19.3 | 18.7 |
| during last 30 | | | | |
| Did this one or more times | 53.4 | 51.3 | 58.7 | 49.4 |

Anal sex within the past 30 days

| | KSU | KSU | KSU | Reference |
|-----------------------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| Never did this activity | 73.6 | 69.6 | 71.5 | 75.2 |
| Have not done this activity | 20.6 | 24.2 | 22.8 | 19.7 |
| during last 30 | | | | |
| Did this one or more times | 5.8 | 6.1 | 5.8 | 5.1 |

Reported means of birth control used among sexually-active college students to prevent pregnancy the last time they had vaginal intercourse

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|--------------------------|-------------|-------------|-------------|------------------|
| Birth control pills | 43.1 | 40.6 | 46.4 | 34.2 |
| Depo Provera | 2.3 | 2.8 | 0.7 | 1.3 |
| Norplant | 0.0 | 0.5 | 0.2 | 0.5 |
| Condoms | 36.8 | 39.8 | 43.8 | 35.8 |
| Diaphragm, cervical cap, | 0.0 | 0.3 | 0.2 | 0.4 |
| sponge | | | | |
| Spermicide | 6.0 | 5.0 | 4.2 | 2.5 |
| Fertility awareness | 3.8 | 2.0 | 2.3 | 2.4 |
| Withdrawal | 20.5 | 17.1 | 23.2 | 15.6 |
| Other method | 5.8 | 5.3 | 5.9 | 6.2 |
| Nothing | 4.7 | 6.0 | 4.7 | 6.3 |

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|-----------------|-------------|-------------|-------------|------------------|
| Usage of Plan B | 8.4 | 9.6 | 11.8 | 11.7 |

G. Nutrition

Within the last 30 days, college students reported:

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|--|-------------|-------------|-------------|------------------|
| Exercising to lose weight | 51.3 | 52.1 | 54.8 | 53.1 |
| Dieting to lose weight | 36.9 | 33.3 | 43.1 | 34.0 |
| Vomiting/Taking laxatives to lose weight | 3.5 | 1.8 | 5.2 | 2.1 |
| Taking diet pills to lose weight | 6.6 | 4.3 | 5.6 | 3.7 |
| Doing none of the above | 41.1 | 42.8 | 33.7 | 38.6 |

College students reported usually eating the following number of servings of fruits and vegetables:

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|---------------------------------|-------------|-------------|-------------|------------------|
| Don't eat fruits and vegetables | 3.7 | 8.5 | 4.5 | 3.7 |
| 1 or 2 per day | 65.2 | 66.7 | 58.8 | 60.8 |
| 3-4 per day | 26.0 | 21.2 | 31.1 | 29.1 |
| 5 or more per day | 5.1 | 3.5 | 5.6 | 6.4 |

College students reported the following behaviors within the past 7 days:

| Participated in vigorous exercise | e for 20 mi | nutes (or 3l |) minutes mod | <i>lerate exercise)</i> |
|-----------------------------------|-------------|--------------|---------------|-------------------------|
| | | | | |

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|----------|-------------|-------------|-------------|------------------|
| 0 days | 38.8 | 34.5 | 30.7 | 28.1 |
| 1-2 days | 29.1 | 32.8 | 30.4 | 32.3 |
| 3-5 days | 26.9 | 28.6 | 30.2 | 32.0 |
| 6+ days | 5.2 | 4.0 | 8.7 | 7.6 |

Exercising to strengthen muscles

| | KSU | KSU | KSU | Reference |
|----------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| 0 days | 45.2 | 45.3 | 42.7 | 42.3 |
| 1-2 days | 29.1 | 29.8 | 27.7 | 28.6 |
| 3-5 days | 20.7 | 22.6 | 25.4 | 24.4 |
| 6+ days | 4.9 | 2.4 | 4.3 | 4.8 |

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|----------|-------------|-------------|-------------|------------------|
| 0 days | 12.6 | 12.7 | 11.3 | 12.1 |
| 1-2 days | 30.5 | 33.4 | 25.2 | 27.8 |
| 3-5 days | 43.8 | 43.0 | 51.1 | 46.2 |
| 6+ days | 13.2 | 10.9 | 12.5 | 13.8 |

Getting enough sleep to feel rested in the morning

Estimated average Body Mass Index (BMI): This figure incorporates reported sex, height, and weight to form a general indicator of physical health:

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|----------------------------|-------------|-------------|-------------|------------------|
| < 18.5 Underweight | 7.2 | 4.8 | 3.1 | 4.8 |
| 18.5 – 24.9 Healthy Weight | 60.3 | 58.8 | 63.8 | 58.6 |
| 25 – 29.9 Overweight | 20.5 | 22.6 | 19.5 | 22.9 |
| 30 – 34.9 Class I Obesity | 8.6 | 8.3 | 7.1 | 8.2 |
| 35 – 39.9 Class II Obesity | 2.5 | 3.1 | 3.8 | 3.4 |
| >= 40 Class III Obesity | 0.9 | 2.4 | 2.6 | 2.1 |

H. Depression

Students reported experiencing the following within the last 12 months:

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------|-------------|-------------|-------------|------------------|
| Never | 5.2 | 6.2 | 5.2 | 8.4 |
| 1-10 times | 64.4 | 64.5 | 64.9 | 64.5 |
| 11+ times | 30.4 | 29.2 | 30.0 | 27.1 |

Feeling overwhelmed by all they had to do

Feeling exhausted (not from physical activity)

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------|-------------|-------------|-------------|---------------------|
| Never | 10.2 | 8.7 | 8.7 | 11.2 |
| 1-10 times | 61.9 | 67.0 | 65.9 | 64.4 |
| 11+ times | 27.8 | 24.3 | 25.4 | 24.3 |

Feeling very sad

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------|-------------|-------------|-------------|------------------|
| Never | 16.8 | 22.3 | 20.5 | 23.5 |
| 1-10 times | 67.9 | 60.8 | 68.2 | 63.5 |
| 11+ times | 15.3 | 16.9 | 11.3 | 13.0 |

Feeling things were hopeless

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------|-------------|-------------|-------------|------------------|
| Never | 29.0 | 33.0 | 29.9 | 38.5 |
| 1-10 times | 59.2 | 53.8 | 60.9 | 51.7 |
| 11+ times | 11.8 | 13.2 | 9.2 | 9.8 |

Feeling so depressed it was difficult to function

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------|-------------|-------------|-------------|------------------|
| Never | 51.0 | 53.4 | 52.2 | 56.8 |
| 1-10 times | 41.0 | 37.1 | 41.8 | 36.2 |
| 11+ times | 8.0 | 9.6 | 5.9 | 7.0 |

Seriously considering attempting suicide

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------|-------------|-------------|-------------|------------------|
| Never | 89.0 | 84.9 | 89.6 | 89.7 |
| 1-10 times | 9.2 | 14.1 | 10.4 | 9.3 |
| 11+ times | 1.8 | 1.0 | 0.0 | 1.0 |

Attempting suicide

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|------------|-------------|-------------|-------------|------------------|
| Never | 97.8 | 98.3 | 98.6 | 98.2 |
| 1-10 times | 1.8 | 1.7 | 1.4 | 1.7 |
| 11+ times | 0.3 | 0.0 | 0.0 | 0.2 |

College students reported diagnosed with depression:

| KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|-------------|-------------|-------------|------------------|
| 16.5 | 16.5 | 16.5 | 16.0 |

| Of those students reporting ever naving been augnosed with depression | | | | |
|---|------|------|------|-----------|
| | KSU | KSU | KSU | Reference |
| | 2003 | 2004 | 2008 | (2007) |
| Diagnosed with depression in | 45.4 | 44.8 | 37.1 | 39.2 |
| last 12 months | | | | |
| Currently in therapy for | 25.0 | 26.0 | 34.3 | 24.2 |
| depression | | | | |
| Currently taking medication for | 44.4 | 36.5 | 43.5 | 35.8 |
| depression | | | | |

Of those students reporting ever having been diagnosed with depression...

Demographics

| Age | | | | |
|---------------|------|------|------|-----------|
| | KSU | KSU | KSU | Reference |
| | 2003 | 2004 | 2008 | (2007) |
| 18-20 years | 47.7 | 47.9 | 40.3 | 53.0 |
| 21 – 24 years | 30.2 | 31.4 | 42.9 | 28.5 |
| 25 – 29 years | 10.1 | 10.8 | 9.2 | 8.1 |
| 30+ years | 12.0 | 9.9 | 7.6 | 10.4 |

Sex

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|--------|-------------|-------------|-------------|------------------|
| Female | 75.5 | 59.4 | 77.8 | 63.3 |
| Male | 24.5 | 40.6 | 20.6 | 33.1 |

Ethnicity

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|----------------------------|-------------|-------------|-------------|---------------------|
| White | 89.0 | 87.5 | 90.2 | 75.2 |
| Black – not Hispanic | 3.8 | 5.5 | 4.2 | 4.7 |
| Hispanic or Latino | 1.1 | 1.0 | 1.4 | 6.9 |
| Asian or Pacific | 2.1 | 3.8 | 2.1 | 9.3 |
| American Indian or Alaskan | 0.5 | 1.2 | 0.7 | 1.6 |
| Native | | | | |
| Other | 2.7 | 2.2 | 2.3 | 3.9 |

Sexual Orientation

| | KSU | KSU | KSU | Reference |
|---------------|------|------|------|-----------|
| | 2003 | 2004 | 2008 | (2007) |
| Heterosexual | 93.1 | 94.1 | 94.0 | 94.0 |
| Gay/Lesbian | 1.1 | 1.5 | 1.9 | 1.6 |
| Bisexual | 4.0 | 2.6 | 2.4 | 2.5 |
| Transgendered | 0.0 | 0.0 | 0.2 | 0.2 |
| Unsure | 1.7 | 1.7 | 1.4 | 1.8 |

Have any kind of health insurance

| | KSU 2003 | KSU 2004 | KSU 2008 | Reference (2007) |
|--------|-------------|-------------|-------------|------------------|
| Yes | 78.8 | 78.5 | 85.1 | 83.8 |
| Unsure | 9.5 | 9.7 | 4.7 | 6.4 |